

SHORT COMMUNICATION**EFFECT OF SOME LICHEN EXTRACTS FROM SRI LANKA ON LARVAE OF *Aedes aegypti* AND THE FUNGUS *Cladosporium cladosporioides***

C. NANAYAKKARA¹, K. BOMBUWELA², S. KATHIRGAMANATHAR², N. K. B. ADIKARAM¹, D. S. A. WIJESUNDARA³, G. N. HARIHARAN⁴, P. WOLSELEY⁵ and V. KARUNARATNE^{2*}

¹ Department of Botany, University of Peradeniya, Peradeniya.

² Department of Chemistry, University of Peradeniya, Peradeniya.

³ Royal Botanic Gardens, Peradeniya.

⁴ M.S. Swaminathan Research Foundation, Chennai, India.

⁵ Department of Botany, The Natural History Museum, U.K.

(Accepted: 26 January 2005)

Abstract: Dichloromethane and methanol extracts of 48 lichens collected from six areas in Sri Lanka were subjected to mosquitolarvicidal assay against the second instar larvae of *Aedes aegypti* and antifungal assay against *Cladosporium cladosporioides*. The following extracts showed significant mosquitolarvicidal activity. CH₂Cl₂: *Cladonia coniocraea*, *Everniastrum* sp., *Parmelina tiliaceae*, *Parmotrema chinense*, *Parmotrema tinctorum*, *Rimelia reticulata*, *Dirinaria applanta*, *Heterodermia leucomelos*, *Roccella montagnei*, *Stereocaulon* sp., *Ocellularia* sp., two species of *Myriotrema*, and *Usnea* sp. MeOH: *Hypogymnia* sp., *H. leucomelos*, *Lepraria atrotomentosa*. The CH₂Cl₂ extracts of *C. coniocraea*, *Hypogymnia* sp., *Lecanora* sp., *Parmeliella* sp., *Everniastrum* sp., *Heterodermia diademata*, *H. leucomelos*, *R. montagnei*, and *Usnea* sp., exhibited potent antifungal activity.

Keywords: *Aedes aegypti*, *Cladosporium cladosporioides*, Lichen extracts, mosquito larvicidal and antifungal activity, Sri Lankan lichens.

INTRODUCTION

Lichens are symbiotic organisms of fungi (mycobionts) and algae (phycobionts) comprising of about 17,000 species recorded worldwide.¹ They commonly grow on rock surfaces, poorly developed soils such as those found in arid lands and boreal-arctic regions and as epiphytes on trees and shrubs.² Lichens synthesize a variety of metabolites with diverse biological activities. For example, lichens possess antibiotic activity, antitumour and antimutagenic activity, anti-HIV activity and plant growth and enzyme inhibitory activity.³ The majority of the compounds responsible for these activities originate from the fungal mycobiont. The general resistance of lichens to insects and microbial attack is attributed to the presence of lichen metabolites.

Lichens have not been subjected to comprehensive bioassays unlike the flowering plants. Recently, however, there have been some reports of screening of lichen metabolites, particularly for antibiotic activity.^{4,5,6,7} Large pharmaceutical companies have initiated screening programs for application of these metabolites and their derivatives in medicine and crop protection.² In this investigation we report the bioactivities of some lichens of Sri Lanka, against the second instar larvae of *Aedes aegypti* and the fungus *Cladosporium cladosporioides*.

METHODOLOGY

Lichen species collected from six selected areas were used in this study. Specimens of lichens were deposited in the National Herbarium at the Royal Botanic Gardens, Peradeniya. Each lichen species (100 g) was washed thoroughly under running water, air-dried and extracted into CH₂Cl₂ (2.5 L) separately in bottle shakers for 24 h at room temperature. The dichloromethane extracts were filtered through a fluted filter paper and the solvents were evaporated under reduced pressure using a rotavapor below 40° C to yield crude extracts. The lichen residues after the extraction into CH₂Cl₂ were extracted into MeOH following the same procedure.

Laboratory cultured larvae of *A. aegypti* were used for this assay.⁸ Five-second instar larvae were introduced into the test solutions (25 ml). Four replicates were used for each experiment. Test solutions were prepared by

* Corresponding author

Table 1: Mosquito larvicidal activity and antifungal activity of the extracts of some lichen species of Sri Lanka

Lichen species	Collection	Mosquito larvicidal activity		Antifungal activity	
		CH ₂ Cl ₂	MeOH	CH ₂ Cl ₂	MeOH
Arthoniaceae					
<i>Arthonia</i> sp. (90) #	PD	-	-	*	*
Cladoniaceae					
<i>Cladonia</i> sp. (31)	PT	1	-	2	-
<i>Cladonia coniocraea</i> (Flörke) Spreng. (55)	HG, HR	3	-	2	-
Collemaaceae					
<i>Collema</i> sp. (02)	HG	-	-	*	*
<i>Leptogium denticulatum</i> * (104)	HG, HR, PD	1	-	-	-
<i>Leptogium saturninum</i> (Dicks.) Nyl. (11)	HG	-	-	*	*
Graphidaceae					
<i>Graphina</i> sp. (100)	PD	1	-	-	-
Parmeliaceae					
<i>Hypogymnia</i> sp. (30)	HG, HR	2	3	3	1
<i>Menegazzia</i> sp. (71)	HG, HR	-	1	*	*
Lecanoraceae					
<i>Lecanora</i> sp. (103)	HG, HR	-	-	*	*
<i>Lecanora</i> sp. (52)	HG, HR	-	1	-	-
<i>Lecanora</i> sp. (80)	HG, HR, PD	2	-	3	1
Letrouitiaceae					
<i>Letrouitia domingensis</i> (Pers.) Haf. & Bellem.(44)	HG, HR, PD	1	-	*	*
Lobariaceae					
<i>Lobaria retigera</i> (Bory) Trevis. (74)	HG	1	1	1	-
<i>Lobaria</i> sp. (73)	HG, HR	1	1	*	*
<i>Pseudocyphellaria argyracea</i> (Bory) Vain. (15)	HG	1	1*	*	
<i>Pseudocyphellaria</i> sp. (76)	HG, HR	2	-	1	-
<i>Sticta ciliaris</i> Mont. et v.d. Bosch (18)	HG	2	-	-	-
Megalosporaceae					
<i>Megalospora</i> sp. (47)	AB, PT, HR	-	1	*	*
Pannariaceae					
<i>Leproloma sipmanianum</i> Kümmerl & Leuckert (79)	BE	1	-	1	-
<i>Pannaria</i> sp. (08)	HG	2	2	1	-
<i>Parmeliella</i> sp. (07)	HG	3	1	2	-
Parmeliaceae					
<i>Everniastrum</i> sp.	HG	3	2	2	1
<i>Hypotrachyna</i> sp. (39)	PT, HG	1	1	1	-
<i>Parmelina tiliacea</i> (Hoffm.) Hale (Elix & McCarthy)	HG, HR	3	2	1	-
<i>Parmotrema chinense</i> (Osbeck) Hale & Ahti	HG, HR	3	1	1	1
<i>Parmotrema crinitum</i> (Ach.) M.Choisy. (24)	HG	2	1	-	-
<i>Parmotrema tinctorum</i> (Nyl.) Hale (72)	HG, HR, PD	3	2	1	-
<i>Rimelia reticulata</i> (63) (Taylor) Hale & A. Fletcher	HG	3	-	-	-
Pertusariaceae					
<i>Pertusaria</i> sp. (77)	BE	1	-	*	*
<i>Pertusaria</i> sp. (105)	HG, PD	2	1	*	*
Physciaceae					
<i>Dirinaria applanta</i> (Fée) Awasthi (78)	HG, HR	3	-	1	-

Lichen Species	Collection	Mosquito larvicidal activity		Antifungal activity	
		CH ₂ Cl ₂	MeOH	CH ₂ Cl ₂	MeOH
<i>Dirinaria confluens</i> (Fr.) Awasthi (99)	HG, HR, PD	2	1	-	1
<i>Heterodermia diademata</i> (Tayl.) Awas. (23)	HR	1	-	2	1
<i>Heterodermia leucomelos</i> (L.) Poelt (46)	HR	3	3	2	1
<i>Heterodermia comosa</i> (Eschw.) Follm. & Redón (4)	HG, HR	1	1	1	-
<i>Pyxine coccifera</i> (Fée) Nyl. (84)	HG, PD	1	1	*	*
Ramalinaceae					
<i>Ramalina</i> sp. (61)	HG, HR	2	1	*	*
Roccellaceae					
<i>Roccella montagnei</i> (Del.) (81)	KU, PD	3	-	2	-
Sphaerophoraceae					
<i>Sphaerophorus</i> sp. (61)	HR	-	-	-	2
Stereocaulaceae					
<i>Stereocaulon</i> sp. (101)	HG	3	-	1	1
Teloschistaceae					
<i>Caloplaca</i> sp. (93)	PD	1	1	*	*
Thelotremataceae					
<i>Ocellularia</i> sp. (09)	HG, HR, PT	3	2	1	-
<i>Myriotrema</i> sp. (48)	HG, HR	3	1	-	-
<i>Myriotrema</i> sp. (14)	HG, HR	3	2	-	-
Trichotheliaceae					
<i>Porina</i> sp. (19)	HG, HR, PT	2	1	*	*
Usneaceae					
<i>Usnea</i> sp. (51)	AB	3	-	3	2
<i>Usnea plicata</i> (L.) (27)	HG, HR	2	-	1	-
(Uncertain)					
<i>Lepraria atrotomentosa</i> Orange & Wolseley (49)	RB	2	3	1	*

AB: Ambewela; BE: Beragala; HG: Hakgala; HR: Horton Plains; KU: Kurunegala; PD: Peradeniya; PT: Pattipola; RB: Ramboda; #: Herbarium specimen number; *: Not studied; -: No activity; For antifungal activity- 1: Low activity (3 or <3mm); 2: Moderate activity (3-20mm); 3: High activity (>20mm). For mosquito larvicidal activity- 1: Low activity (1-40 %); 2: Moderate activity (41-90 %); 3: High activity (91-100 %).

dissolving the extract (100 mg) in a minimum volume (< 1 ml) of acetone and diluting with water up to 200 ml (500 ppm). Polyethyleneglycol (PEG) (120 µL) was added to the mixture to homogenise the test solution. A series of concentrations were prepared by diluting the 500 ppm solution with water. After the introduction of larvae, a small portion of powdered fish food was added to each beaker and the beakers were covered with watch glasses. In all experiments an untreated control, which contained all the ingredients except the lichen extract, was used. The number of moribund or dead larvae was recorded after 24 and 48 h.

Antifungal assays were carried out on extracts by dissolving the extract (2 mg) in either

CH₂Cl₂ or MeOH (1 ml). Benlate (methyl-1-(butylcarbonyl)-2-benzamidizolecarbamate), a commercial fungicide (1 mg, 50% active ingredient), was also dissolved in water (1 ml) and used in the assays for comparison with the extracts.⁹

TLC plates (silica gel 60 PF₂₅₄₊₃₆₆, 0.5 mm ×20 cm ×20 cm) were spotted with extracts at about one inch from the base. The plates were developed in CH₂Cl₂ or 5% MeOH in CH₂Cl₂ and then air-dried (overnight) to evaporate all traces of remaining solvents. A spore suspension of *Cladosporium cladosporioides* in Czapek-dox nutrient solution (CNS) was sprayed onto the plates and the plates were incubated in a moist

chamber at 28 ± 2 °C for two days. On every plate run, Benlate was spotted as the standard. Inhibition zones appeared white against a background of green mycelia. The diameter of zones in which the growth was inhibited, which were approximately circular, was measured.

RESULTS

Dichloromethane and methanol extracts of 48 species of lichens collected from eight locations in Sri Lanka were subjected to mosquito larvicidal assay against the second instar larvae of *Aedes aegypti* and antifungal assay against *Cladosporium cladosporioides*. The CH₂Cl₂ extracts of *Cladonia coniocraea*, *Hypogymnia* sp., *Parmeliella* sp., *Everniastrum* sp., *Parmelina tiliaceae*, *Parmotrema chinense*, *Parmotrema tinctorum*, *Rimelia reticulata*, *Dirinaria applanta*, *Heterodermia leucomelos*, *Roccella montagnei*, *Stereocaulon* sp., *Ocellularia* sp., two species of *Myriotrema* and *Usnea* sp., and MeOH extracts of *Hypogymnia* sp., *H. leucomelos*, *Lepraria atrotomentosa* showed highly significant mosquito larvicidal activity. The CH₂Cl₂ extracts of *C. coniocraea*, *Hypogymnia* sp., *Lecanora* sp., *Parmeliella* sp., *Everniastrum* sp., *Heterodermia diademata*, *H. leucomelos*, *R. montagnei*, and *Usnea* sp., exhibited potent antifungal activity. *C. coniocraea*, *Hypogymnia* sp., *Everniastrum* sp., *Parmeliella* sp., *Lecanora* sp., *H. leucomelos*, *R. montagnei* and *Usnea* sp., showed both mosquito larvicidal and antifungal activity.

Acknowledgement

Authors acknowledge financial support from NSF, NRC and IFS, Sweden.

References

1. Huneck S. (1999). The significance of lichens and their metabolites. *Naturwissenschaften* **86**: 559-570.
2. Crittenden P. D. & Porter N. (1991). Lichen-forming fungi: Potential sources of novel metabolites. *Focus* **9**: 409.
3. Huneck S. & Yoshimura I. (1996). *Identification of lichen substances*. Springer, Berlin Heidelberg New York.
4. Calder V. L., Cole A. L. & Walker J. R. L. (1986). Antibiotic compounds from New Zealand plants III. A survey of some New Zealand plants for antibiotic substances. *Journal of the Royal Society New Zealand* **16**: 169.
5. Cosar G., Tumbay E., Zeybek N. & Ozer A. (1998). The antibacterial and antifungal effect of some lichens growing in Turkey I. *Evernia prunastri*, *Pseudevernia furfuracea* and *Alectoria capillaries*. *Fitoterapia* **59**: 505.
6. Ingolfsdottir K., Bloomfield S. F. & Hylands P. J. (1985). In vitro evaluation of the antimicrobial activity of lichen metabolites as potential preservatives. *Antimicrobial Agents and Chemotherapy* **28**: 289-292.
7. Lawrey J. D. (1986). Biological role of lichen substances. *Bryologist* **89**: 111.
8. Bandara K. A. N. P., Kumar V., Jacobsson U. & Molleyres P. (2000). An insecticidal piperidine alkaloid from *Microcos paniculata* stem bark. *Phytochemistry* **54**: 29.
9. Klarman W. L. & Stanford J. B. (1968). Isolation and purification of antifungal principles from infected soy beans. *Life Sciences* **7**: 1095.