

Evaluation of the heat resistance of pathogenic *Listeria monocytogenes* in milk and milk products in Sri Lanka

VS Jayamanne¹ and U Samarajeewa²

¹Department of Food Science & Technology, Faculty of Agriculture, University of Ruhuna, Kamburupitiya, Sri Lanka

²Department of Food Science & Technology, Faculty of Agriculture, University of Peradeniya, Peradeniya, Sri Lanka

Accepted: 26th July 2010

ABSTRACT

Listeria monocytogenes, which is a food-borne pathogen often isolated from milk and milk products, causes listeriosis in pregnant women, newborns, older adults and immuno-suppressed people. *Listeria* is said to be more resistant to heat treatments and the present study investigated the effects of current heat treatments on inactivation of *Listeria* in milk. Two isolates of *L. monocytogenes*, namely FSTLC2 and FSTLC55, were introduced to sterilized milk at population levels of 10^2 and 10^7 cfu/ml and were subjected to various temperature-time combinations. *Listeria* was enumerated using the FDA *Listeria* Enrichment Broth (LEB; Oxoid Ltd) and Modified Oxford Agar (MOX; Oxoid Ltd) medium. The Thermal Death Time (TDT) and Thermal Death Rate (TDR) curves were also constructed using thermal resistance data. The slope of the TDT curve (z) and the slope of the TDR curve (D ; Decimal Reduction Time) were also determined using the statistical software package Minitab (release 8.21). It was observed that the normal pasteurization treatments of 62.8°C for 30 min (LTLT; Low Temperature Long Time) and 71.7°C for 15sec (HTST; High Temperature Short Time) appear to be adequate to destroy a *Listeria* population of 10^2 CFU/ml in milk, but not a population of 10^7 cfu/ml. A z value of 7.6°C and a $D_{71.7}^{\circ\text{C}}$ value of 2.9sec were observed for *L. monocytogenes* in milk. The current heat treatment techniques used in the dairy industry need to be reviewed and necessary modifications introduced to ensure a complete elimination of this pathogen. Moreover, the z and $D_{71.7}^{\circ\text{C}}$ values observed in the present study may be used in devising sound heat treatment techniques as well as suitable Hazard Analysis Critical Control Points (HACCP) systems in the dairy industry.

Keywords: *Listeria monocytogenes*, Thermal inactivation, Heat resistance, Pasteurization, Thermal Death Time, Thermal Death Rate

INTRODUCTION

Listeria monocytogenes, which is a Gram-positive, facultatively anaerobic, cold tolerant, salt tolerant, non-spore forming and non-acid fast rod often found in soil and water, and faeces of animals, is a relatively recent food-borne pathogen reported mostly from developed countries (Maijala *et al.* 2001; Karakolev 2009). *L. monocytogenes* is a food-borne pathogen of high concern to the food industry as its ubiquitous occurrence in the environment can lead to contamination of foods. *L. monocytogenes* has been isolated from soil and water, fresh vegetables, potato, faeces of animals, cattle and faeces of humans, newborn infants, silage, raw beef and chicken, and Soymilk. The toxic substance produced during the exponential growth phase of *L. monocytogenes* is designated listeriolysin O (hemolysin), which causes hemolysis of blood agar in *in-vitro* studies. When *L. monocytogenes* is contracted through the oral route, it apparently colonizes the intestinal tract by mechanisms that are poorly understood. From the intestinal tract, the organism

invades tissues including the placenta in pregnant women, and enters the blood stream, from which it reaches other susceptible body tissues. Listeriosis, the illness caused by this bacterium, affects mostly pregnant women, elderly, newborns, and the immuno-suppressed adults due to AIDS, alcoholism, diabetes, cancer, cardiovascular disease, kidney disease, renal transplant, and corticosteroid therapy (Rocourt *et al.* 2000). When susceptible adults contract the disease, meningitis and septicemia are the most commonly recognized symptoms. The non-pregnant healthy individuals who are not immuno-suppressed are fairly resistant to infection by *L. monocytogenes*. Infected pregnant women may experience only mild flu-like illness, and the illness can be transmitted from mother to the fetus through the placenta resulting in abortion, premature birth, stillbirth, or serious health problems for the newborn child. The mortality rate of the disease is approximately 25% (USDA 1999).

Milk and milk products have received much attention because they have been reported to harbor *L. monocytogenes* to a much higher extent compared

*Corresponding author; vsjayamanne@fst.ruh.ac.lk

Paper presented at the 2nd National Symposium, Faculty of Agriculture, University of Ruhuna

to other foods. *L. monocytogenes* has been isolated from cow milk, goat milk, cheese, and pasteurized milk in Sri Lanka (Jayamanne and Samarajeewa 2001). There have been reports of *Listeria* outbreaks associated with dairy products (Farber and Peterkin 1991). Cow milk has been implicated in food-borne fatal listeriosis. The ability of *L. monocytogenes* strains to proliferate in raw milk even under refrigerated condition is well documented. Hitherto only two suspected human cases of listeriosis have been reported in Sri Lanka. The first is two children suffering from meningitis (Jayasundara *et al.* 1962; Withana and Mirando 1967), and two cases (children) with headache, fever and a rash (Wijesundera *et al.* 1992). *L. monocytogenes* has been isolated from goslings, chickens, buffaloes, and goats in Sri Lanka (Bandaranayaka and Nair 1962; Bandaranayaka and Fernando 1962).

Though milk and milk products were reported to transmit food-borne listeriosis, only two studies have been reported in Sri Lanka to ascertain the safety of the dairy products. In a survey on *Listeria* in foods, it was demonstrated that the percentage of *L. monocytogenes* positive samples was highest among vegetables (49%), second in chicken (34%), and lowest in dairy products (26%) (Gunaseena *et al.* 1995). They further observed that 5 pasteurized milk (31%), 3 raw milk (25%) and 4 ice cream (33%) samples contained *L. monocytogenes*. In a similar study, of the samples tested (265), 39 samples (15%) contained virulent *L. monocytogenes*. Cow milk (29%), goat milk (27%), pasteurized milk (17%), and cheese (33%) samples contained virulent strains of *L. monocytogenes* (Jayamanne & Samarajeewa 2001).

Listeria is said to be more heat resistant to heat processing than other foods, especially when buffered by proteinaceous foods. Although there have been reports on the presence and thermal properties of *Listeria* in milk and milk products in other countries, no such studies have been reported from Sri Lanka. Therefore, the present study investigated the thermal resistance of isolates of *L. monocytogenes* in milk and milk products in Sri Lanka.

MATERIALS AND METHODS

Separate experiments were carried out to determine the optimum pasteurization-time combination required to inactivate *Listeria* in milk, to determine the thermal resistance of *Listeria* from pasteurized milk and to determine the z and $D_{71.7}^0$ values of *Listeria* in bovine milk during heat processing.

Determination of the optimum pasteurization temperature-time combination required to inactivate *Listeria*

A commercial brand of sterilized milk (fat ~3.5%, protein ~4%) available in the market was selected as the milk based food system for the experiment. Sterilized milk was first tested for *Listeria* using the FDA *Listeria* Enrichment Broth (LEB; Oxoid Ltd., Basingstoke, UK) and Modified Oxford Agar medium (MOX; Oxoid Ltd.) in order to ensure that milk was not contaminated with *Listeria*.

In the enumeration process of the organism, *Listeria* in milk was first enriched in FDA *Listeria* Enrichment Broth. Samples (25 ml) were blended with LEB (225ml) at 12,000rpm for 2min in a Waring blender. The resulting solution was incubated at 35°C for 48h for selective enrichment and the selective isolation of *Listeria* was done in MOX. Enriched samples were next streaked onto MOX Agar plates and incubated at 35°C and examined for colonies with black halos after 24, 48 and 72h. Single colonies growing on the MOX medium were isolated and transferred to MOX agar slants in McCartney bottles and incubated (35°C) until there was a sufficient growth. The MOX agar slants were stored in the refrigerator (3°C) pending confirmatory morphological and biochemical tests. MOX agar slants were sub-cultured at 3-month interval to ensure the viability of organisms. The tests [microscopy on wet mount, Gram staining, catalase test, methyl red (MR) test, Vogus-Proskauer (VP) test, H₂S production, color changes in Triple Sugar Iron Agar, color changes in Urea Agar, hydrolysis of aesculin and CAMP test] were carried out to confirm the presence of *Listeria* in the food samples.

L. monocytogenes FSTLC2 (University of Peradeniya Culture Collection), isolated previously in one of our previous studies (Jayamanne and Samarajeewa, 2001) from raw cow milk was enriched in LEB at 35°C for 48h. A loopful of enriched *Listeria* culture was aseptically transferred to McCartney bottles containing 20ml of Tryptose Broth (Oxoid Ltd.), incubated for another 24h at 35°C. After 24h of incubation in Tryptose Broth, the *Listeria* cells at the late stationary phase were harvested by centrifugation at 6000 × g for 20min. *Listeria* was introduced to sterilized milk to have an approximate initial *Listeria* population of 10²cfu/ml and the other with 10⁷cfu/ml. Two experiments were carried out separately, one with an approximate initial *Listeria* population of 10²cfu/ml and the other with 10⁷cfu/ml.

Sterilized milk with *Listeria* was subjected to different temperature regimes (from 50 to 80°C at 5°C increments) for different time periods (5, 10, 15, 20, 25 and 30sec, and 1, 2.5, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 110, 120,

130, 140, and 150min) on a constant temperature water bath (Kayagaki-DC3). Predetermined come-up times were allowed prior to the actual experiments. Samples (1ml) in duplicate were aseptically taken in the beginning, and after heat treatments, and enriched in LEB for 48h at 35°C. Enriched cultures were streaked on to MOX Agar plates and incubated at 35°C for 48h, and examined for typical *L. monocytogenes* colonies with black/brown halos. The minimum time periods, required for the total inactivation of *L. monocytogenes* with regard to different temperature regimes, were recorded.

Determination of the thermal resistance of *L. monocytogenes* isolates from pasteurized Milk

L. monocytogenes FSTLC55 (University of Peradeniya Culture Collection) previously from pasteurized milk was aseptically introduced to sterilized milk (at populations of 10^2 cfu/ml) and was subjected to different temperature regimes and time periods as described earlier. After giving heat treatments, samples were enriched in LEB and inoculated on MOX Agar plates and examined for any sign of the survival of *Listeria*.

Determination of z and D values for *L. monocytogenes* in bovine milk

The Thermal Death Time (TDT) curve was constructed using previous data, i.e., inactivation time for *Listeria* with regard to different temperature values used. The slope of the TDT curve (z) was determined using the statistical package Minitab (release 8.21). The decimal reduction time (D) values for *L. monocytogenes* in cow milk with regard to HTST pasteurization treatment (71.7°C for 15 sec) was determined in a separate experiment. The commercial brand of sterilized milk used in the previous studies was used in this experiment as well. *L. monocytogenes* FSTLC2 from the stored culture collection was enriched first in LEB for 48 h at 35°C, transferred to Tryptose Broth, and incubated for further 24h at 35°C. *Listeria* cells at the late stationary phase were harvested at $6,000 \times g$ for 20min and introduced to sterilized milk (250ml) in an Erlenmeyer flask to make a final *Listeria* population of $\sim 10^5$ cfu/ml. The Erlenmeyer flask was incubated at 71.7°C on a constant temperature water bath. Samples (1ml) in duplicate were aseptically taken from Erlenmeyer flask at 3 sec intervals. These samples were enriched in LEB for 48 h at 35°C. A dilution series (from 1:10 to 1:100000) with buffered water was prepared, and two MOX Agar plates from each dilution were inoculated using the surface plate count technique. MOX Agar plates were incubated at 35°C for 48h and a surface count was taken using a digital colony counter (Kayagaki-DC3). Logarithmic of the survival number of or-

ganisms (Y axis) was plotted against time (X axis). The time required to reduce the number of organisms by one log cycle is the decimal reduction time (D) in relation to the temperature regime used, 71.7°C used in this experiment.

RESULTS AND DISCUSSION

It was observed in the present study that the normal pasteurization protocols such as HTSTb and LTLT treatments are adequate to inactivate a low *Listeria* population of 10^2 cfu/ml but not a high population of 10^7 cfu/ml.

Evaluation of the most effective pasteurization temperature-time combination

The observed temperature-time combinations required to ensure complete inactivation of the *L. monocytogenes* FSTLC2 from raw cow milk under laboratory conditions is given in Table 1. It was observed that the normal pasteurization treatments of 62.8 °C for 30 min (LTLT), and 71.7 °C for 15 sec (HTST) appear to be adequate to destroy a *Listeria* population of 10^2 cfu/ml in milk, but not at population of 10^7 cfu/ml as shown in Table 1, Fig. 1a and Fig. 1b. It was reported by many early scientists that LTLT (Mackey and Bratchell, 1989) and HTST (Bunning *et al.* 1992) treatments are adequate to destroy *Listeria* in milk. Numerous conflicting reports concerning the unusual heat resistance of *L. monocytogenes* in milk can be found in the early literature. *L. monocytogenes* survived 15sec in milk at 100°C (Ozgen, 1952). *L. monocytogenes* was not completely inactivated until the milk was held at 65 °C for 5 min, 75 °C for 2 min, or 80°C for 3-5 min (Stenberg and Hammainen, 1955). *L. monocytogenes* survived 30-40sec at 65°C, 10 sec at 75°C, and ~1 sec at 85°C (Dedie and Schulze 1957). But, later scientists found many inadequacies in the methodologies employed in these early experiments, and the validity of these

Table 1 Thermal inactivation times of *L. monocytogenes* (FSTLC2) suspended in milk at populations of 10^2 and 10^7 cfu/ml

Temperature (°C)	Thermal inactivation time	
	10^2 cfu/ml	10^7 cfu/ml
50	140min	160min
55	90min	120min
60	30min	40min
65	15min	20min
70	15sec	25sec
75	10sec	15sec
80	5sec	10sec




Figure 1a: Comparison of LTLT treatment and temperature-time combinations in eliminating *L. monocytogenes* FSTLC2 in milk at populations of 10^2 and 10^7 fu/ml




Fig. 1b: Comparison of HTST treatment and temperature-time combinations in eliminating *L. monocytogenes* FSTLC2 in milk at populations of 10^2 and 10^7 cfu/ml

findings remains controversial.

The temperature-time combinations required for inactivation of *L. monocytogenes* suspended in milk at population levels of 10^2 and 10^7 cfu/ml as observed in the model experiment (Table 1) are compared with LTLT and HTST pasteurization treatments (Figure 1a and 1b). The figures indicate that the risk area (representing temperature-time combinations of treatments) increases with the initial population of *Listeria* in milk. Thus, for a high *Listeria* population level of 10^7 cfu/ml, the LTLT and HTST pasteurization treatments fall inside the risk area.

It is safe to conclude that for a low *Listeria* population of less than 10^2 cfu/ml, LTLT and HTST treatments may always be inside the safe area. When the *Listeria* population increases, the chances of LTLT and HTST treatments falling in the boundary or risk area also increases, thus, making pasteurization ineffective. It is therefore evident that, if milk is contaminated with a heavy population of *Listeria* ($>10^7$ cfu/ml), neither LTLT nor HTST treatments could be considered as reliable heat processing techniques. It was observed in a comparable study that the complete inactivation of *L. monocytogenes* in milk with an initial population of 10^6 - 10^7 *Listeria* cfu/ml cannot be accomplished within 30 min at 62, 72, 82, or even 92°C (Donnelly *et al.* 1986). The thermal inactivation of *Listeria* in milk is associated with the initial load of the organism, the fat content, heat adaptation of the organism, and state of the organism in milk (free suspension or internalized).

In Sri Lanka, milk products such as curd, yogurt, and cottage cheese are produced at home level and in micro-enterprises. The heat processing techniques they use in eliminating microorganisms may not always be carefully controlled standard methods. They boil milk without taking into consideration the time factor. Even though such a boiling is adequate to eliminate a low *Listeria* population of

less than 10^2 cfu/ml, it may not be adequate to eliminate a higher initial *Listeria* population of 10^7 cfu/ml. Even though the possibility of the presence of a *Listeria* population of 10^7 cfu/ml in raw milk was observed to be very low in this study, the observations cannot be extended to the variety of conditions operating during milking, collection, transport, storage and processing of milk leaving much room for rapid increase of *Listeria* populations to an infectious dose. The ubiquitous nature and the ability of *L. monocytogenes* to survive harsh environmental conditions also should not be ignored. The absence of proper chilling facilities, and long time-gap in between milking and processing add to the severity of the problem. With regard to pasteurized milk, there is always an associated risk, especially if proper thermal inactivation conditions are not practiced, because pasteurized milk is a ready-to-consume product, which would not be further processed.

Thermal resistance of *L. monocytogenes* FSTLC55 isolated from pasteurized milk

The present study showed that the thermal inactivation time for the *L. monocytogenes* FSTLC55, an isolate which had been obtained from a commercial brand of pasteurized milk in Sri Lanka, is higher than that for an isolate from raw cow milk (FSTLC2), at the same population of 10^2 cfu/ml (Table 2).

The thermal inactivation times for *L. monocytogenes* isolate from raw cow milk (FSTLC2), introduced at different initial populations of 10^2 and 10^7 cfu/ml, and thermal inactivation times for the isolate from pasteurized milk (FSTLC55), at population of 10^2 cfu/ml, are illustrated in Figure 2. At the same population of 10^2 cfu/ml, *L. monocytogenes* isolate from pasteurized milk (FSTLC55) showed higher thermal resistance properties than

Table 2: Comparison of thermal inactivation times for *Listeria* isolates from pasteurized milk (FSTLC55) and raw cow milk (FSTLC2) at population of 10^2 cfu/ml


Temperature (°C)	Thermal inactivation time	
	Isolate from raw cow milk	Isolate from pasteurized milk
50	140min	160min
55	90min	130min
60	30min	45min
65	15min	30min
70	15sec	50sec
75	10sec	25sec
80	5sec	15sec

the isolate from raw cow milk (FSTLC2) (Figure 2). It is also evident that the isolate from raw cow milk (FSTLC2), when suspended at 10^7 cfu/ml, showed higher thermal resistance than when suspended at populations 10^2 cfu/ml (Figure 2). It is known that when the *Listeria* population increases in milk, the required thermal inactivation time also increases. The notable feature here is that the isolate from pasteurized milk needed higher heat treatment for inactivation than a population of 10^7 cfu/ml from raw milk. The elevated thermal resistance of *L. monocytogenes* isolates, observed especially in *L. monocytogenes* isolate from pasteurized milk, could be associated with following reasons:


It was reported that the fat in milk protects *L. monocytogenes* from heat. *L. monocytogenes* cells surrounded by milk fat are relatively much more heat stable (MacDonald and Sutherland 1993). The protective effect of milk fat is important during the maximum death region of the survival curve. Milk foods should not be processed at temperatures below 60°C because the shoulder regions become more pronounced and thermal processing may be under-estimated (Chhabra 1999). This is due to the buffering action and the poor thermal conductivity exerted by fat in milk.

In cases of naturally acquired listerial mastitis, the pathogen is normally not freely suspended in milk but rather exists as a facultative intracellular bacterium within phagocytic leukocytes (neutrophils and macrophages) present in milk (Tilney and Portnoy 1989). The partly intracellular nature of *L. monocytogenes* makes it possible that cells of the pathogen inside leukocytes could be partially protected from thermal inactivation and thus are more able to survive pasteurization than freely suspended cells of the bacterium in milk.

If *L. monocytogenes* is shed from infected cows, which have developed fever, the pathogen may have grown at elevated temperatures. It was reported that when *Listeria* was grown at 43°C before heating and recovered by anaerobic incubation after

**Figure 2: Comparison of thermal inactivation times for *L. monocytogenes* isolates from pasteurized milk (FSTLC55) and raw cow milk (FSTLC2)**

the treatment, the $D_{62.8\text{ }^\circ\text{C}}$ value was 243 sec as compared with 36 sec when the pathogen was grown at 37°C and plated aerobically after the heat inactivation (Knabel *et al.*, 1990) suggesting that cows who have developed fever may harbor organisms which are more heat resistant. It is also believed that *Listeria* cells inside macrophages are exposed to anaerobic conditions. It is possible that a few heat-injured cells remaining after the minimum HTST milk pasteurization may grow under anaerobic conditions that may exist in phagocytes. It is reported that adaptation to the environmental stresses such as high/low temperature, acidic and oxidative conditions, and starvation occurs in all bacteria, including *L. monocytogenes* (Mims 1987). An increasing number of parameters such as temperature, salt, pH, heat shock, and oxidative stress, have been shown to affect virulence gene expression in *L. monocytogenes* (Brehm *et al.* 1996). Resistance of *L. monocytogenes* to heat or other lethal factors can be increased by heat shock or adaptation to other stresses. It was reported that bacteria respond to heat shock by synthesizing new proteins, termed heat-shocked proteins (HSPs), and these proteins usually increase the thermo-tolerance of microorganisms (Craig *et al.* 1993). Heat-shock and other environmental stresses are also reported to affect the virulence of *L. monocytogenes*. Environmental stresses are sensed by pathogens as sig-

**Figure 3: Regressed TDT curve for *L. monocytogenes* FSTLC2 suspended in bovine milk at populations 10^5 cfu/ml**

nals for expression of virulence factors to enhance survival (Archer, 1996). Heat shock may also increase the virulence of *L. monocytogenes* enhancing more listeriolysin O production.

It is evident that *L. monocytogenes*, at times, shows unexpectedly higher thermal resistance to normal pasteurization protocols practiced in the dairy processing industry. Therefore, it is essential that the thermal processing techniques be reviewed at regular intervals in order to ensure the effectiveness of them.

Determination of z value for *L. monocytogenes* using the Thermal Death Time (TDT) curve

A z value of 7.6°C was observed for *L. monocytogenes* FSTLC2 isolated from raw cow milk. The z value is the temperature in centigrade required to make a tenfold change in the Thermal Death Time (TDT) curve. The regression line of the TDT curve for the *L. monocytogenes* FSTLC2 suspended in sterilized milk is illustrated in Figure 3. Thermal processing is the most widely used method to preserve food and to check harmful microorganisms to ensure safety for human consumption. TDT and TDR curves give general insight into the thermal inactivation of an organism during heat processing. The z values of 6.5 (Bradshaw *et al.* 1987), 7.3 (Bunning *et al.* 1986), 8.0 (Bunning *et al.* 1988), and 6.3 (Bradshaw *et al.* 1985) were reported for *L. monocytogenes* in milk. The apparent difference of the z value of the present study from other reported studies could be attributed to the food system used in the experiments. Some scientists used skim milk (Bradshaw *et al.* 1987), whereas other scientists used whole milk (Bunning *et al.* 1988). The fat content is higher in whole milk thus giving protective effects to *L. monocytogenes* against heat processing. This is partly due to the poor thermal conductivity in high fat milk foods. The higher the fat content in milk foods, the higher will be the z value. Food system used in the present study was sterilized whole milk. Therefore, it is possible that the relatively high z value (7.6°C) observed is due to the buffering action of fat in milk. The z value observed in the present study is of the same order as reported z values, but high.

Determination of the Decimal Reduction Time (D) for *L. monocytogenes* using the TDR curve

A decimal reduction time ($D_{71.7^{\circ}\text{C}}$) of 2.9 sec was observed for *L. monocytogenes* FSTLC2- which had been isolated from raw cow milk- suspended in sterilized whole milk at populations 10^5cfu/ml (Fig. 4). Thermal Death Rate (TDR) curve was constructed using the survival number of *L. monocytogenes* isolate suspended in sterilized milk with processing time (sec) at 71.7°C . By definition, D value




Figure 4: Regressed TDR curve for *L. monocytogenes* FSTLC2 suspended in bovine milk at populations 10^5cfu/ml at 71.7°C

Predicted regression model: $X=16.5-2.84Y$

X =Processing time (sec), Y =log survival number of *Listeria*, $r^2=91.0$, $p=0.003$

is the time required to eliminate 90% of the organism at a given temperature, which is 71.7°C in the present experiment.

Some of the reported $D_{71.7^{\circ}\text{C}}$ values in milk for *L. monocytogenes* include 1.7sec (Bradshaw *et al.* 1987), 3.1 and 5.0sec (Bunning *et al.* 1988), and 1.9 sec (Bunning *et al.* 1986). The differences, in the observed D value in the present study and reported D values, are once again due to the different milk food systems used in the respective experiments. Some scientists used skim milk (Bradshaw *et al.* 1987), whereas other scientists used whole milk (Bunning *et al.* 1988) in thermal resistance studies. Sterile whole milk was used in the present study as well. The fat content is higher in whole milk thus giving some protective effects to *L. monocytogenes* against heat processing.

The higher the D and z values, the longer it takes to eliminate *L. monocytogenes* from milk foods. It is apparent that it takes longer times to eliminate *Listeria* added to sterile whole milk where fat content is high. The observed $D_{71.7^{\circ}\text{C}}$ value of the present study is comparable with those of other thermal resistance studies, and lies in between reported extreme $D_{71.7^{\circ}\text{C}}$ values of 0.9sec (Bradshaw *et al.* 1985) and 5.0sec (Bunning *et al.* 1988). These observations suggest that *L. monocytogenes* isolates used in the present study show average thermal resistance properties.

The z and D values observed could be effective tools for food processing in Hazard Analysis and Critical Control Point (HACCP) plans, risk assessment models, and product development formulations in Sri Lanka perspective. Given the consequences of human listeriosis outbreaks, it may be necessary to encourage milk processing plants in general and milk pasteurizing plants in particular in Sri Lanka to review their thermal processing techniques to ascertain whether they are on par with the observed z and D values.

CONCLUSIONS

Normal pasteurization techniques such as LTLT and HTST protocols can effectively eliminate a low population level (10^2 CFU/ml) of *Listeria*, but not a high population (10^7 CFU/ml) level in milk. Therefore, the current heat treatment techniques used in the dairy industry need to be reviewed and necessary modifications introduced in order to ensure a complete elimination of this pathogen in milk and milk products in Sri Lanka. Moreover, the z (7.6°C) and $D_{71.7^\circ\text{C}}$ (2.9sec) values observed in the present study may be used in devising sound heat treatment techniques as well as suitable Hazard Analysis Critical Control Points (HACCP) systems in the dairy industry.

Acknowledgements

The authors gratefully acknowledge the assistance given by Mr PK Liyanage (Dept of Food Science & Technology, University of Peradeniya) and Mrs S Abeysekare (Dept of Food Science & Technology, University of Ruhuna) in preparing microbiological media and taking readings.

REFERENCES

- Archer DL 1996 Preservation microbiology and safety: evidence that stress enhances virulence and triggers adaptive mutations. *Trends Food Sci. Technol.* 7: 91-95.
- Bandaranayaka A and Nair VK 1962 Listeriosis in buffaloes. *Cey. Vet. J.* 10: 42-45.
- Bandaranayaka A, and Fernando WWHS 1962 A note on the isolation of *Listeria monocytogenes* from the heart blood of a goat. *Cey. Vet. J.* 10: 46-47.
- Bradshaw JG, Peeler JT, Corwin JJ, Hunt JM and Twedt RM 1987 Thermal resistance of *Listeria monocytogenes* in dairy products. *J. Food Prot.* 50: 543-544.
- Bradshaw, J.G., Peeler, J.T., Corwin, J.J., Hunt, J.M., Tierney, J.T., Larkin, E.P., and Twedt, R.M. (1985). Thermal resistance of *Listeria monocytogenes* in milk. *J. Food Protect.* 48: 743-755.
- Brehm, K., Kreft, J., Ripio, M.T., and Vazquez-Boland, J.A. (1996). Regulation of virulence gene expression in pathogenic *Listeria*. *Microbiol. Sem.* 12: 219-236.
- Bunning, V.K., Crawford, R.G., Bradshaw, J.G., Peeler, J.T., Tierney, J.T., and Twedt, R.M., (1986). Thermal resistance of intracellular *Listeria monocytogenes* cells suspended in raw bovine milk. *Appl. Environ. Microbiol.* 52: 1398-1402.
- Bunning VK, Donnelly CW, Peeler JT, Briggs EH, Bradshaw JG, Crawford RG, Beliveau CM and Tierney JT 1988 Thermal inactivation of *Listeria monocytogenes* within bovine milk phagocytes. *Appl. Environ. Microbiol.* 54: 364-370.
- Bunning VK, Crawford RG, Tierney JT and Peeler JT 1992 Thermotolerance of heat-shocked *Listeria monocytogenes* in milk exposed to high-temperature short-time pasteurization. *Appl. Environ. Microbiol.* 58(6): 2096-2098.
- Chhabra AT, Carter WH, Linton RH and Cousin MA 1999 A predictive model to determine the effects of pH, milkfat, and temperature on thermal inactivation of *Listeria monocytogenes*. *J. Food Prot.* 62(10): 1143-1149.
- Dieck K and Schulze D 1957 Die Hitzeresistenz von *Listeria monocytogenes* in Milch. *Berl. Munchener tierarztl. Wochenscher.* 70: 231-232.
- Donnelly CW, Briggs EH and Baigent GJ 1986. Analysis of raw milk for the epidemic serotype of *Listeria monocytogenes* linked to an outbreak of listeriosis in California. *J. Food Prot.* 49: 846-847 (Abstract).
- Farber JM and Peterkin PI 1991 *Listeria monocytogenes*, a food-borne pathogen. *Microbiol. Rev.* 55: 476-511.
- Fleming DW, Cochi SL, Macdonald KL, Brondum, J, Hayes PS, Plikaytis BD, Holmes MB, Audurier A, Broome CV and Reingold AL 1985 Pasteurized milk as a vehicle of infection in an outbreak of listeriosis. *New Eng. J. Med.* 312(7): 404-407.
- Gunasena DK, Kodikara CP, Ganepola K, Widanapathirana S 1995 Occurrence of *Listeria monocytogenes* in food in Sri Lanka. *J. Natn. Sci. Coun. Sri Lanka.* 23(3): 107-114.
- Jayamanne VS, and Samarajeewa U 2001 Incidence and detection of *Listeria monocytogenes* in milk and milk products of Sri Lanka. *Trop. Agric. Res.* 13: 42-50.
- Jayasundera LBT, Goonathilleke PDP and Fernando NI 1962 A fatal case of meningitis due to *Listeria (Erysipelothrix) monocytogenes*. *Cey. Med. J.* 7: 221-225.
- Knabel SJ, Walker HW, Hartman PA and Mendonca AF 1990 Effects of growth temperature and strictly anaerobic recovery on survival of *Listeria monocytogenes* during pasteurization. *Appl. Environ. Microbiol.* 56: 370-376.
- Karakolev R 2009 Incidence of *Listeria monocytogenes* in beef, pork, raw-dried and raw-smoked sausages in Bulgaria. *Food Control.* 20(10): 953-955.
- Mackey BM and Bratchell N 1989 The heat resistance of *Listeria monocytogenes*. *Let. Appl. Microbiol.* 9: 89-94.
- MacDonald F and Sutherland AD 1993 Effect of heat treatment on *Listeria monocytogenes* and Gram-negative bacteria in sheep, cow, and goat milks. *J. Appl. Bacteriol.* 75(4): 336-343.
- Maijala R, Lyytikainen O, Johansson T, Autio T,

- Aalto T, Haavisto L and Honkanen-Buzalski T 2001 Exposure of *Listeria monocytogenes* within an epidemic caused by butter in Finland. *Int. J. Food Microbiol.* 70:97-109.
- Mims CA 1987 The encounter of the microbe with the phagocytic cell. *In: The Pathogenesis of Infectious Diseases.* Orlando, FL: Academic Press. Pp. 63-91.
- Ozgen H 1952 Zur Serologic der *Listeria monocytogenes*. *Z. Tropenmed.* 4: 40-45.
- Rocourt J, Jacquet C and Reilly A 2000 Epidemiology of human listeriosis and seafoods. *Int. J. Food Microbiol.* 67:197-209.
- Stenberg H and Hammainen T 1955 On determination *in vitro* of the resistance of *Listeria monocytogenes* to sodium chloride and heat on experimental monocytosis in albino mice. *Nord. Vet. Med.* 7: 853-868.
- Tilney LG and Portnoy DA 1989 Actin filaments and the growth, movement, and spread of the intracellular bacterial parasite, *Listeria monocytogenes*. *J. Cell Biol.* 109: 1597-1608.
- United States Department of Agriculture 1999 Consumer Information from USDA, Food Safety Education Office, Food Safety and inspection service, USDA, Washington D.C. 20250-3700, USA.
- Wijesundera C de S, Pinto MRM and Navaratnam C 1992 Two cases of listeriosis. *Cey. Med. J.* 37: 21-23.
- Withana KA and Mirando EH 1967 Meningitis caused by *Listeria monocytogenes*. *Cey. Med. J.* 12: 37-39.